

# LOCAL RINGS OF BOUNDED COHEN–MACAULAY TYPE

GRAHAM J. LEUSCHKE AND ROGER WIEGAND

ABSTRACT. Let  $(R, \mathfrak{m}, k)$  be a local Cohen–Macaulay (CM) ring of dimension one. It is known that  $R$  has finite CM type if and only if  $R$  is reduced and has bounded CM type. Here we study the one-dimensional rings of bounded but infinite CM type. We will classify these rings up to analytic isomorphism (under the additional hypothesis that the ring contains an infinite field). In the first section we deal with the complete case, and in the second we show that bounded CM type ascends to and descends from the completion. In the third section we study ascent and descent in higher dimensions and prove a Brauer–Thrall theorem for excellent rings.

Let  $R$  be a Cohen–Macaulay (CM for short) local ring. We say that  $R$  has finite, respectively bounded, CM type provided there are only finitely many indecomposable maximal Cohen–Macaulay (MCM)  $R$ -modules up to isomorphism, respectively, there is a bound on the multiplicities of the indecomposable MCM  $R$ -modules. The one-dimensional CM local rings of finite CM type have been completely classified [DR], [GR], [Ç1], [Ç2], [ÇWW]. One consequence of the characterization is that a one-dimensional CM local ring  $R$  has finite CM type if and only if the completion  $\widehat{R}$  has bounded CM type and is reduced.

Here we study the one-dimensional CM local rings of bounded but infinite CM type. In [LW2] we classified the complete equicharacteristic hypersurface singularities of dimension one having bounded but infinite CM type: Up to isomorphism, the only ones are  $k[[X, Y]]/(Y^2)$  and  $k[[X, Y]]/(XY^2)$ . In §1 of this paper we consider complete equicharacteristic rings that are not hypersurfaces, and we show that only one additional isomorphism type arises (Theorem 1.5). In the second section we prove that bounded CM type ascends to and descends from the completion. Our main results in dimension one are summarized in Theorem 2.4. In the third section of the paper we study ascent and descent in higher dimensions.

## 1. ONE-DIMENSIONAL COMPLETE LOCAL RINGS

We begin by quoting two results we will need from [LW2]. Recall that an  $R$ -module  $M$  is said to *have (constant) rank  $r$*  [SS] provided  $K \otimes_R M \cong K^r$ , where  $K$  is the total quotient ring of  $R$  (obtained by inverting all nonzerodivisors). We denote by  $\nu_R(M)$  the number of generators required for  $M$  as an  $R$ -module.

---

*Date:* April 15, 2003 (filename: BCMTrevised1.tex).

Leuschke’s research was supported by an NSF Postdoctoral Fellowship, and Wiegand’s was supported by grants from the NSA and the NSF.

**Proposition 1.1.** [LW2, Theorem 3.1] *Let  $R$  be a complete, equicharacteristic hypersurface of dimension one (i.e.,  $R \cong k[[X, Y]]/(f)$  for some field  $k$  and some non-zero non-unit  $f$  in the formal power series ring  $k[[X, Y]]$ ). Then  $R$  has bounded but infinite CM type if and only if either  $R \cong k[[X, Y]]/(Y^2)$  or  $R \cong k[[X, Y]]/(XY^2)$ . Further, if  $R$  has unbounded CM type, then  $R$  has, for each positive integer  $r$ , an indecomposable MCM module of constant rank  $r$ .*

**Proposition 1.2.** *Let  $(R, \mathfrak{m})$  be a one-dimensional local CM ring. Assume that either*

- (1)  *$R$  has multiplicity at least 4, or*
- (2)  *$R$  has a birational extension  $S$  such that  $\nu_R(S) = 3$  and  $\nu_R(\frac{\mathfrak{m}S}{\mathfrak{m}}) > 1$ .*

*Then  $R$  has, for each positive integer  $r$ , an indecomposable MCM module of constant rank  $r$ .*

While Proposition 1.2 is not stated explicitly in [LW2], it follows immediately from Lemma 2.2 and Theorem 2.3 there. Also, one needs (0.5) and (2.1) of [ÇWW] to deduce the “Further” statement in Proposition 1.1. Before stating our main result (Theorem 1.5), we state two lemmas that will be useful here and in the next section. Part (1) of the first lemma is due to Bass [B, (7.2)]. We include the proof here, since the context in [B] is a bit different from ours.

**Lemma 1.3.** *Let  $(R, \mathfrak{m})$  be a Gorenstein local ring of dimension one with total quotient ring  $K$ . Let  $E = \text{End}_R(\mathfrak{m}) = \{\lambda \in K \mid \lambda\mathfrak{m} \subseteq \mathfrak{m}\}$ . Assume that  $E$  is local.*

- (1) *If  $M$  is an indecomposable MCM  $R$ -module and  $M \not\cong R$ , then  $M$  is an indecomposable MCM  $E$ -module (naturally, that is, in such a way that the  $R$ -module structure induced by the natural map  $R \hookrightarrow E$  agrees with the original structure).*
- (2) *Every indecomposable MCM  $E$ -module is an indecomposable MCM  $R$ -module.*

*In particular,  $R$  has finite (respectively, bounded) CM type if and only if  $E$  has finite (respectively, bounded) CM type.*

*Proof.* (1) (Bass, [B]) Since there is no surjection from  $M$  to  $R$ , we have  $M^* = \text{Hom}_R(M, \mathfrak{m})$ , which is an  $E$ -module. Therefore  $M \cong M^{**}$  is an  $E$ -module as well. Clearly  ${}_E M$  is indecomposable and (since  $E$  is local) maximal Cohen–Macaulay.

(2) Since  $R \hookrightarrow E$  is module-finite, any MCM  $E$ -module  $N$  is MCM as an  $R$ -module. Since  $R \hookrightarrow E$  is birational and  ${}_R N$  is torsion-free, any  $R$ -endomorphism of  $N$  is automatically  $E$ -linear. It follows that  $N$  is indecomposable as an  $R$ -module.

The final statement is immediate from (1) and (2). □

We are indebted to Tom Marley (private communication) for showing us the following lemma, and to the anonymous referee for suggesting an improvement to the proof. Let  $e(R)$  denote the multiplicity of the local ring  $R$ .

**Lemma 1.4.** *Let  $(R, \mathfrak{m}, k)$  be a one-dimensional local CM ring with  $k$  infinite, and suppose  $e(R) = \nu_R(\mathfrak{m}) = 3$ . Let  $N$  be the nilradical of  $R$ . Then:*

- (1)  $N^2 = 0$ .

- (2)  $\nu_R(N) \leq 2$ .
- (3) If  $\nu_R(N) = 2$ , then  $\mathfrak{m}$  is generated by three elements  $x, y, z$  such that  $\mathfrak{m}^2 = \mathfrak{m}x$  and  $N = Ry + Rz$ .
- (4) If  $\nu_R(N) = 1$ , then  $\mathfrak{m}$  is generated by three elements  $x, y, z$  such that  $\mathfrak{m}^2 = \mathfrak{m}x$ ,  $N = Rz$ , and  $yz = z^2 = 0$ .

*Proof (Marley).* We note that  $R$  has minimal multiplicity [Ab] and hence  $\mathfrak{m}$  has reduction number 1. Since the residue field is infinite, then, there is an element  $x \in \mathfrak{m}$  such that  $\mathfrak{m}^2 = x\mathfrak{m}$ . Since  $R$  is CM,  $x$  is a nonzerodivisor. We recall the formula [S, (1.1)]

$$(1.4.1) \quad \nu_R(J) \leq e(R) - e(R/J)$$

for an ideal  $J$  of height 0 in a one-dimensional CM local ring  $R$ . Now the image of  $x$  is a reduction element for  $R/N^2$ , so the right-hand side of (1.4.1), with  $J = N^2$ , is  $e(R/(x)) - e(R/(Rx + N^2))$ . But  $N^2 \subseteq \mathfrak{m}^2 \subset Rx$ , so this expression is 0. We conclude from (1.4.1) that  $N^2 = 0$ . Similarly, putting  $J = N$ , we see that  $\nu_R(N) \leq 2$ .

For (3) and (4), we observe that  $N/\mathfrak{m}N \subseteq \mathfrak{m}/\mathfrak{m}^2$ , that is, minimal generators of  $N$  are also minimal generators of  $\mathfrak{m}/\mathfrak{m}^2$  and of  $\mathfrak{m}$ . Indeed, since  $x$  is a nonzerodivisor mod  $N$ ,  $N \cap x\mathfrak{m} = xN$ ; it follows that  $N \cap \mathfrak{m}^2 = \mathfrak{m}N$ , and the map  $N/\mathfrak{m}N \rightarrow \mathfrak{m}/\mathfrak{m}^2$  is injective. This proves (3).

To complete the proof of (4), let  $z$  generate  $N$ . Choose any  $w$  such that  $\mathfrak{m} = Rx + Rw + Rz$ . Now  $zw = xg$  for some  $g$ , and (as  $x$  is a nonzerodivisor)  $g = bz$  for some  $b \in R$ . Then  $z(w - bx) = 0$ , and we may take  $y = w - bx$ .  $\square$

**Theorem 1.5.** *Let  $k$  be an infinite field. The following is a complete list, up to  $k$ -isomorphism, of the one-dimensional, complete, equicharacteristic, CM local rings with bounded but infinite CM type and with residue field  $k$ :*

- (1)  $k[[X, Y]]/(Y^2)$ ;
- (2)  $T := k[[X, Y]]/(XY^2)$ ;
- (3)  $E := \text{End}_T(\mathfrak{m}_T)$ , where  $\mathfrak{m}_T$  is the maximal ideal of  $T$ . We have a presentation  $E \cong k[[X, Y, Z]]/(XY, YZ, Z^2)$ .

Moreover, if  $(R, \mathfrak{m}, k)$  is a one-dimensional, complete, equicharacteristic CM local ring and  $R$  does not have bounded CM type, then  $R$  has, for each positive integer  $r$ , an indecomposable MCM module of constant rank  $r$ .

*Proof.* The rings in (1) and (2) have bounded but infinite CM type by Proposition 1.1. To show that  $E$  has bounded CM type, it suffices, by Lemma 1.3, to check that  $E$  is local, a fact that will emerge in the next paragraph, where we verify the presentation of  $E$  given in (3). We routinely use decapitalization to denote specialization of variables.

The element  $x + y$  is a nonzerodivisor of  $T$ , and the fraction  $z := \frac{y^2}{x+y}$  is easily checked to be in  $E := \text{End}_T(\mathfrak{m}_T)$  but not in  $T$ . Now  $E = \text{Hom}_T(\mathfrak{m}_T, T)$  since  $\mathfrak{m}_T$  does not have  $T$  as a direct summand, and it follows by duality over the Gorenstein ring  $T$  that  $E/T \cong \text{Ext}_R^1(T/\mathfrak{m}_T, T) \cong k$ . Therefore  $E = T[z]$ . Since  $z^2 = \frac{y^2(x+y)^2}{(x+y)^2} \in \mathfrak{m}_T$ ,  $E$  is local. One verifies the relations  $xz = 0, yz = y^2 = z^2$  in  $E$ . Thus the map  $S := k[[X', Y', Z']] \rightarrow E$  (sending

$X' \mapsto x, Y' \mapsto y$ , and  $Z' \mapsto z$ ) induces a surjection from  $A := S/(X'Z', Y'^2 - Y'Z', Y'Z' - Z'^2)$  to  $E$ . We will build an inverse map, but first we note that the change of variables  $X' = X, Y' = Y + Z, Z' = Y$  transforms  $A$  to the ring  $B := k[[X, Y, Z]]/(XY, YZ, Z^2)$ . Since  $B/(z)$  is one-dimensional and  $z^2 = 0$ ,  $B$  and  $A$  are one-dimensional.

To build the inverse map, we note that we have  $x'y^2 = 0$  in  $A$ , so we have, at least, a map from  $T$  to  $A$  taking  $x$  to  $x'$  and  $y$  to  $y'$ . Next, we note that the defining ideal  $I$  of  $A$  is the ideal of  $2 \times 2$  minors of the matrix  $\varphi = \begin{bmatrix} X' & Y' \\ Y' - Z' & 0 \\ 0 & Z' \end{bmatrix}$ . By (the converse of) the Hilbert–Burch theorem [BH, (1.4.16)],  $I$  has a free resolution

$$0 \longrightarrow S^2 \xrightarrow{\varphi} S^3 \longrightarrow I \longrightarrow 0.$$

Therefore  $A$  has projective dimension 2 over  $S$  and hence depth 1. Therefore  $A$  is Cohen–Macaulay, and, since  $\mathfrak{m}_A^2 = (x' + y')\mathfrak{m}$ , it follows that  $x' + y'$  is a non-zerodivisor of  $A$ . Now our map  $T \rightarrow A$  extends to a map  $T[\frac{1}{x+y}] \rightarrow A[\frac{1}{x'+y'}]$ , and the restriction of this map provides the desired map from  $E$  to  $A$ .

We now know that each of the rings on our list has bounded but infinite CM type. To show that the list is complete and to prove the “Moreover” statement, assume now that  $(R, \mathfrak{m}, k)$  is a one-dimensional, complete, equicharacteristic CM local ring with  $k$  infinite and having infinite CM type. Suppose, moreover, that  $R$  does *not* have indecomposable MCM modules of arbitrarily large (constant) rank. We will show that  $R$  is isomorphic to one of the rings on the list.

If  $R$  is a hypersurface, Proposition 1.1 tells us that  $R$  is isomorphic to either  $k[[X, Y]]/(Y^2)$  or  $k[[X, Y]]/(XY^2)$ . Thus we assume that  $\nu_R(\mathfrak{m}) \geq 3$ . But  $e(R) \leq 3$  by Proposition 1.2. Therefore we may assume that  $e(R) = \nu_R(\mathfrak{m}) = 3$ . Thus we are in the situation of Lemma 1.4. Moreover, the nilradical  $N$  of  $R$  is non-trivial, by [CWW, (0.5), (1.2)].

We claim that  $N$  is principal. If not, then by (3) of Lemma 1.4, we can find elements  $x, y, z$  in  $R$  such that

$$(1.5.1) \quad \mathfrak{m} = Rx + Ry + Rz, \quad \mathfrak{m}^2 = \mathfrak{m}x, \quad \text{and} \quad N = Ry + Rz$$

Put  $S := R[\frac{y}{x^2}, \frac{z}{x^2}] = R + R\frac{y}{x^2} + R\frac{z}{x^2}$ , and note that  $\mathfrak{m}S = \mathfrak{m} + R\frac{y}{x} + R\frac{z}{x}$ . It is easy to verify (by clearing denominators) that  $\{1, \frac{y}{x^2}, \frac{z}{x^2}\}$  is a minimal generating set for  $S$  as an  $R$ -module, and that the images of  $\frac{y}{x}$  and  $\frac{z}{x}$  form a minimal generating set for  $\frac{\mathfrak{m}S}{\mathfrak{m}}$ . Thus we are in case (2) of Proposition 1.2, and our basic assumption is violated. This proves our claim that  $N$  is principal.

Using Lemma 1.4(4), we find elements  $x, y, z$  in  $R$  such that

$$(1.5.2) \quad \mathfrak{m} = Rx + Ry + Rz, \quad \mathfrak{m}^2 = \mathfrak{m}x, \quad N = Rz, \quad \text{and} \quad yz = z^2 = 0.$$

Since  $y^2 \in \mathfrak{m}x \subset Rx$ , we see that  $R/Rx$  is a three-dimensional  $k$ -algebra. Further, since  $\cap_n(Rx^n) = 0$ , it follows that  $R$  is finitely generated (and free) as a module over the discrete valuation ring  $V := k[[x]]$ .

We claim that  $R = V + Vy + Vz$  (and therefore  $\{1, y, z\}$  is a basis for  $R$  as a  $V$ -module). To see this, we note that  $R = V[[y, z]] = V[[y]] + Vz$ , since  $yz = z^2 = 0$ . Let  $h \in V[[y]]$ , say,  $h = v_0 + v_1y + v_2y^2 + \dots$ , with  $v_i \in V$ . For each  $n \geq 1$  write  $y^{n+1} = xh_n$ , with  $h_n \in \mathfrak{m}^n$ .

(This is possible since  $\mathfrak{m}^{n+1} = x\mathfrak{m}^n$ .) Then  $v_2y^2 + v_3y^3 + \dots = x(v_2h_1 + v_3h_2 + \dots) \in xR$ . Therefore  $h \in V + Vy + xR$ , and it follows that  $R = V + Vy + Vz + xR$ . Our claim now follows from Nakayama's Lemma.

In order to understand the structure of  $R$  we must analyze the equation that puts  $y^2$  into  $x\mathfrak{m}$ . Thus we write  $y^2 = x^r q$ , where  $r \geq 1$  and  $q \in \mathfrak{m} - \mathfrak{m}^2$ . Write  $q = \alpha x + \beta y + \gamma z$ , with  $\alpha, \beta, \gamma \in V$ . Since  $x$  is a non-zerodivisor and  $yz = z^2 = 0$ , we see immediately that  $\alpha = 0$ . Thus we have

$$y^2 = x^r(\beta y + \gamma z)$$

with  $\beta, \gamma \in V$ ; moreover, at least one of  $\beta, \gamma$  must be a unit of  $V$  (since  $q \notin \mathfrak{m}^2$ ).

We claim that  $r = 1$ . For suppose  $r \geq 2$ . Put  $v := \frac{y}{x^2}$  and  $w := \frac{z}{x^2}$ . The relations  $vw = w^2 = 0$  and  $v^2 = x^{r-2}(\beta v + \gamma w)$  show that  $S := R[v, w]$  is a module-finite birational extension of  $R$ . Moreover,  $\nu_R(S) = 3$  and it is easy to see that  $\frac{\mathfrak{m}S}{\mathfrak{m}}$  is minimally generated by the images of  $\frac{y}{x}$  and  $\frac{z}{x}$ . The desired contradiction now follows from Proposition 1.2.

Thus we have

$$(1.5.3) \quad y^2 = x(\beta y + \gamma z)$$

with  $\beta, \gamma \in V$ , and at least one of  $\beta, \gamma$  is a unit of  $V$ . We will produce a hypersurface subring  $A := V[[g]]$  of  $R$  such that  $R = \text{End}_A(\mathfrak{m}_A)$ . We will then show that  $A \cong k[[X, Y]]/(XY^2)$ , and the proof will be complete.

**Case 1:**  $\beta$  is a unit. Consider the subring  $A := V[y] \subset R$ . From (1.5.3) we see that  $xz \in A$ , and it follows easily that  $z \in \text{End}_A(\mathfrak{m}_A)$ , so  $R \subset \text{End}_A(\mathfrak{m}_A)$ . As before (since  $A$  is Gorenstein)  $R = \text{End}_A(\mathfrak{m}_A)$ .

**Case 2**  $\beta$  is not a unit (whence  $\gamma$  is a unit). This time we put  $A := V[y + z] \subset R$ . The equation

$$xy(1 - \beta\gamma^{-1}) = x(y + z) - \gamma^{-1}(y + z)^2$$

shows that  $xy \in A$ . Therefore  $xz \in A$  as well, and as before we conclude that  $R = \text{End}_A(\mathfrak{m}_A)$ .

By Lemma 1.3,  $A$  has infinite CM type but does not have indecomposable MCM modules of arbitrarily large constant rank. Moreover,  $A$  cannot have multiplicity 2, since it has a module-finite birational extension of multiplicity greater than 2. By Proposition 1.1,  $A \cong k[[X, Y]]/(XY^2)$ , as desired.  $\square$

## 2. ASCENT AND DESCENT IN DIMENSION ONE

In this section we show that bounded CM type passes to and from the  $\mathfrak{m}$ -adic completion of an equicharacteristic one-dimensional CM local ring  $(R, \mathfrak{m}, k)$  with  $k$  infinite. Contrary to the situation in higher dimension (see Theorem 3.3 below), we need not assume that  $R$  is excellent with an isolated singularity. Indeed, in dimension one this assumption would make  $\widehat{R}$  reduced, in which case finite and bounded CM type are equivalent [CWW]. We do, however, insist that  $k$  be infinite, in order to use the crucial fact from §1 that failure of bounded CM type implies the existence of MCM modules of unbounded *constant* rank and also to use the explicit equations worked out in [BGS] for the indecomposable MCM modules over  $T := k[[X, Y]]/(XY^2)$ .

Given a local ring  $R$  with completion  $\widehat{R}$  and a finitely generated module  $M$  over  $\widehat{R}$ , we say  $M$  is *extended* (from  $R$ ) provided there is a finitely generated  $R$ -module  $A$  such that  $\widehat{A} \cong M$ . The following proposition appears, in a narrower context, in the 2002 University of Nebraska Ph.D. thesis of M. Arnavut [Ar]. The argument is adapted from [We, (1.5)].

**Proposition 2.1.** *Let  $R$  be a one-dimensional CM local ring with completion  $\widehat{R}$ , and let  $K$  be the total quotient ring of  $\widehat{R}$ . Let  $M$  and  $N$  be finitely generated  $\widehat{R}$ -modules such that  $K \otimes_{\widehat{R}} M \cong K \otimes_{\widehat{R}} N$ . Then  $M$  is extended if and only if  $N$  is extended.*

*Proof.* Assume  $N$  is extended, say,  $N \cong \widehat{A}$ , where  ${}_R A$  is finitely generated. Choose an  $R$ -module homomorphism  $\varphi : M \rightarrow N$  such that  $\varphi \otimes 1_K$  is an isomorphism. We obtain an exact sequence

$$0 \rightarrow V \rightarrow M \xrightarrow{\varphi} N \xrightarrow{\pi} W \rightarrow 0,$$

in which both  $V$  and  $W$  are torsion modules (therefore of finite length). Let  $L = \varphi(M) = \text{Ker}(\pi)$ . Now  $\text{Hom}_{\widehat{R}}(N, W) = \text{Hom}_{\widehat{R}}(\widehat{A}, \widehat{W}) = (\text{Hom}_R(A, W))^\wedge = \text{Hom}_R(A, W)$ , since  $W$  and  $\text{Hom}_R(A, W)$  have finite length. Therefore there is a homomorphism  $\rho \in \text{Hom}_R(A, W)$  such that  $\widehat{\rho} = \pi$ . Letting  $B = \text{Ker}(\rho)$ , we see that  $\widehat{B} \cong L$ .

Next, we consider the short exact sequence

$$(2.1.1) \quad 0 \rightarrow V \rightarrow M \rightarrow L \rightarrow 0,$$

viewed as an element of  $\text{Ext}_{\widehat{R}}^1(L, V) = \text{Ext}_R^1(B, V)$  (again, because  $V$  and  $\text{Ext}_R^1(B, V)$  have finite length). Therefore (2.1.1) is the completion of a short exact sequence

$$0 \rightarrow V \rightarrow C \rightarrow B \rightarrow 0.$$

Then  $\widehat{C} \cong M$ , as desired.  $\square$

A finitely generated module  $M$  over a Noetherian ring  $R$  is said to be *generically free* provided  $M_P$  is  $R_P$ -free for each  $P \in \text{Ass}(R)$ . For a generically free  $R$ -module  $M$ , we let  $\text{rank}_P(M)$  denote the rank of the free  $R_P$ -module  $M_P$ , for  $P \in \text{Ass}(R)$ .

**Corollary 2.2.** *Let  $(R, \mathfrak{m})$  be a one-dimensional CM local ring with completion  $\widehat{R}$ , and let  $M$  be a generically free  $\widehat{R}$ -module. Then  $M$  is extended from  $R$  if and only if  $\text{rank}_P(M) = \text{rank}_Q(M)$  whenever  $P$  and  $Q$  are minimal primes of  $\widehat{R}$  lying over the same prime of  $R$ . In particular, every  $\widehat{R}$ -module of constant rank is extended from an  $R$ -module (necessarily of the same constant rank).*

*Proof.* Suppose  $M \cong \widehat{W}$ , and let  $P$  and  $Q$  be primes of  $\widehat{R}$  lying over  $p \in \text{Spec}(R)$ . Let  $r = \text{rank}_P(M)$ . We have a flat local homomorphism  $R_p \rightarrow \widehat{R}_P$ . It follows from faithfully flat descent [EGA, (2.5.8)] that  $W_p$  is  $R_p$ -free of rank  $r$ . From the change of rings  $R_p \rightarrow \widehat{R}_Q$  we see that  $\text{rank}_Q(M) = r$ . This proves the “only if” implication and the parenthetical remark in the last sentence of the statement.

For the converse, let  $\{p_1, \dots, p_s\}$  be the minimal primes of  $R$ , and let  $r_i = \text{rank}_P(M_P)$  for  $P$  in the fiber over  $p_i$ . Let  $J_1 \cap \dots \cap J_s$  be a primary decomposition of  $(0)$  in  $R$ , with  $\sqrt{J_i} = p_i$ .

(Since  $R$  is CM,  $\mathfrak{m} \notin \text{Ass}(R)$ .) Put  $W = \bigoplus_{i=1}^s (R/J_i)^{r_i}$ . Then  $K \otimes_{\widehat{R}} \widehat{W} \cong K \otimes_R M$ . By Proposition 2.1,  $M$  is extended.  $\square$

Here is our main result of this section.

**Theorem 2.3.** *Let  $(R, \mathfrak{m}, k)$  be a one-dimensional equicharacteristic CM local ring with completion  $\widehat{R}$ . Assume that  $k$  is infinite. Then  $R$  has bounded CM type if and only if  $\widehat{R}$  has bounded CM type. If  $R$  has unbounded CM type, then  $R$  has, for each  $r$ , an indecomposable MCM module of constant rank  $r$ .*

*Proof.* Assume that  $\widehat{R}$  does not have bounded CM type. Fix a positive integer  $r$ . By Theorem 1.5 we know that  $\widehat{R}$  has an indecomposable MCM module  $M$  of constant rank  $r$ . By Corollary 2.2 there is a finitely generated  $R$ -module  $N$ , necessarily MCM and with constant rank  $r$ , such that  $\widehat{N} \cong M$ . Obviously  $N$  too must be indecomposable.

Assume from now on that  $\widehat{R}$  has bounded CM type. If  $\widehat{R}$  has *finite* CM type, the same holds for  $R$ , [ÇWW]. Therefore we assume that  $\widehat{R}$  has infinite CM type. Then  $\widehat{R}$  is isomorphic to one of the rings of Theorem 1.5:  $k[[X, Y]]/(Y^2)$ ,  $T := k[[X, Y]]/(XY^2)$ , or  $E := \text{End}_T(\mathfrak{m}_T)$ . If  $\widehat{R} \cong k[[X, Y]]/(Y^2)$ , then  $e(R) = 2$ , and  $R$  has bounded CM type by [LW2, (2.1)]. Suppose for the moment that we have verified bounded CM type for any local ring  $S$  whose completion is isomorphic to  $E$ . If, now,  $\widehat{R} \cong T$ , put  $S := \text{End}_R(\mathfrak{m})$ . Then  $\widehat{S} \cong E$ , whence  $S$  has bounded CM type. Therefore so has  $R$ , by Lemma 1.3.

Therefore we may assume that  $\widehat{R} = E$ . Our plan is to examine each of the indecomposable non-free  $E$ -modules and then use Proposition 2.1 and Corollary 2.2 to determine exactly which MCM  $E$ -modules are extended from  $R$ . From now on we use the presentation for  $E = \widehat{R}$  given in the proof of Theorem 1.5:  $E \cong k[[X, Y, Z]]/(XZ, Y^2 - YZ, YZ - Z^2)$ . By Lemma 1.3 the indecomposable non-free MCM  $E$ -modules are exactly the indecomposable non-free MCM  $T$ -modules, namely, the cokernels of the following matrices over  $T$  (see [BGS, (4.2)]):

$$(2.3.1) \quad [y]; \quad [xy]; \quad [x]; \quad [y^2]$$

$$(2.3.2) \quad \alpha := \begin{bmatrix} y & x^k \\ 0 & -y \end{bmatrix}; \quad \beta := \begin{bmatrix} xy & x^{k+1} \\ 0 & -xy \end{bmatrix}; \quad \gamma := \begin{bmatrix} xy & x^k \\ 0 & -y \end{bmatrix}; \quad \delta := \begin{bmatrix} y & x^{k+1} \\ 0 & -xy \end{bmatrix}.$$

Let  $P := (x)$  and  $Q := (y)$  be the two minimal prime ideals of  $T$ . Note that  $T_P \cong k((Y))$  and  $T_Q \cong k((X))[Y]/(Y^2)$ . With the exception of  $U := \text{Coker}[y]$  and  $V := \text{Coker}[xy]$ , each of the modules in (2.3.1) and (2.3.2) is generically free, with  $\varphi$ ,  $(\text{rank}_P \text{Coker}(\varphi), \text{rank}_Q \text{Coker}(\varphi))$  given in the following list:

$$(2.3.3) \quad [x], (1, 0); \quad [y^2], (0, 1); \quad \alpha, (0, 1); \quad \beta, (2, 1); \quad \gamma, (1, 1); \quad \delta, (1, 1).$$

Let  $M$  be a MCM  $\widehat{R}$ -module, and write

$$(2.3.4) \quad M \cong (\oplus_{i=1}^a A_i) \oplus (\oplus_{j=1}^b B_j) \oplus (\oplus_{k=1}^c C_k) \oplus (\oplus_{l=1}^d D_l) \oplus U^e \oplus V^f,$$

where the  $A_i, B_j, C_k, D_l$  are indecomposable generically free modules of ranks  $(1, 0)$ ,  $(0, 1)$ ,  $(1, 1)$ ,  $(2, 1)$  (and, again,  $U = \text{Coker}([y])$  and  $V = \text{Coker}([xy])$ ).

Suppose first that  $R$  is a domain. Then  $M$  is extended if and only if  $b = a + d$  and  $e = f = 0$ . Now the indecomposable MCM  $R$ -modules are those whose completions have  $(a, b, c, d, e, f)$  minimal and non-trivial with respect to these relations. (We are using implicitly the fact (see [Wi2, (1.2)] for example) that for two finitely generated  $R$ -modules  $N_1$  and  $N_2$ ,  $N_1$  is isomorphic to a direct summand of  $N_2$  if and only if  $\widehat{N}_1$  is isomorphic to a direct summand of  $\widehat{N}_2$ .) The only possibilities are  $(0, 0, 1, 0, 0, 0)$ ,  $(1, 1, 0, 0, 0, 0)$  and  $(0, 1, 0, 1, 0, 0)$ , and we conclude that the indecomposable  $R$ -modules have rank 1 or 2.

Next, suppose that  $R$  is reduced but not a domain. Then  $R$  has exactly two minimal prime ideals, and we see from Corollary 2.2 that every generically free  $\widehat{R}$ -module is extended from  $R$ ; however, neither  $U$  nor  $V$  can be a direct summand of an extended module. In this case, the indecomposable MCM  $R$ -modules are generically free, with ranks  $(1, 0)$ ,  $(0, 1)$ ,  $(1, 1)$  and  $(2, 1)$  at the minimal prime ideals.

Finally, we assume that  $R$  is not reduced. We must now consider the two modules  $U$  and  $V$  that are not generically free. We will see that  $U := \text{Coker}[y]$  is always extended and that  $V := \text{Coker}[xy]$  is extended if and only if  $R$  has two minimal prime ideals. Note that  $U \cong Txy = Exy$  (the nilradical of  $E = \widehat{R}$ ), and  $V \cong Ty = Ey$ .

The nilradical  $N$  of  $R$  is of course contained in the nilradical  $Exy$  of  $\widehat{R}$ . Moreover, since  $Exy \cong E/(0 :_E xy) = E/(y, z)$  is a faithful cyclic module over  $E/(y, z) \cong k[[x]]$ , every non-zero submodule of  $Exy$  is isomorphic to  $Exy$ . In particular,  $N\widehat{R} \cong Exy$ . This shows that  $U$  is extended.

Next we deal with  $V$ . The kernel of the map  $Ey \rightarrow Exy$  (multiplication by  $x$ ) is  $Ey^2$ . Thus we have a short exact sequence

$$(2.3.5) \quad 0 \longrightarrow Ey^2 \longrightarrow V \longrightarrow U \longrightarrow 0.$$

Now  $Ey^2 = Ty^2 = \text{Coker}([x])$  is generically free of rank  $(1, 0)$ , and since the total quotient ring  $K$  (of both  $T$  and  $\widehat{R}$ ) is Gorenstein we see that

$$KV \cong Ky^2 \oplus KU.$$

If, now,  $R$  has two minimal primes, every generically free  $\widehat{R}$ -module is extended, by Corollary 2.2. In particular,  $Ey^2$  is extended, and by Proposition 2.1 so is  $V$ . Thus every indecomposable MCM  $\widehat{R}$ -module is extended, and  $R$  has bounded CM type.

If, on the other hand,  $R$  has just one minimal prime ideal, then the module  $M$  in (2.3.4) is extended if and only if  $b = a + d + f$ . The  $\widehat{R}$ -modules corresponding to indecomposable MCM  $R$ -modules are therefore  $U, V \oplus W$ , where  $W$  is some generically free module of rank  $(0, 1)$ , and the modules of constant rank 1 and 2 described above.  $\square$

We conclude this section with a summary of the main results of §§1 and 2.



**Theorem 2.4.** *Let  $(R, \mathfrak{m}, k)$  be an equicharacteristic one-dimensional local CM ring with  $k$  infinite. Then  $R$  has bounded but infinite CM type if and only if the completion  $\widehat{R}$  is isomorphic to one of the following:*

- (1)  $k[[X, Y]]/(Y^2)$ ;
- (2)  $T := k[[X, Y]]/(XY^2)$ ;
- (3)  $E := \text{End}_T(\mathfrak{m}_T)$ , where  $\mathfrak{m}_T$  is the maximal ideal of  $T$ .

Moreover, if  $R$  does not have bounded CM type, then  $R$  has, for each positive integer  $r$ , an indecomposable MCM module of constant rank  $r$ .

### 3. ASCENT, DESCENT, AND BRAUER–THRALL IN HIGHER DIMENSIONS

In this section we study ascent and descent of bounded CM type to and from the completion in dimension greater than one. We prove that bounded CM type ascends to the completion of an excellent CM local ring with an isolated singularity. An easy corollary of this result is a generalization of the Brauer–Thrall theorem of Yoshino and Dieterich [Y, (6.4)]. See Theorem 3.4.

For descent, we have a less complete picture. We show that bounded CM type descends from the completion of a Henselian local ring, and we investigate the case of a two-dimensional normal local domain such that the completion is also a normal domain. Example 3.5 indicates why descent is less tractable than ascent.

For Henselian rings, ascent and descent are easy:

**Proposition 3.1.** *Let  $R$  be a Henselian local ring with completion  $\widehat{R}$ . If  $\widehat{R}$  has bounded CM type, then so has  $R$ . Conversely, if  $R$  has bounded CM type and  $\widehat{R}$  has at most an isolated singularity, then  $\widehat{R}$  has bounded CM type.*

*Proof.* Assume  $\widehat{R}$  has bounded CM type, and let  $M$  be an indecomposable MCM  $R$ -module. Since  $R$  is Henselian, the endomorphism ring  $E := \text{End}_R(M)$  is local, meaning  $E/J$  is a division ring (where  $J$  is the Jacobson radical of  $E$ ). Passing to  $\widehat{R}$ , we observe that  $\text{End}_{\widehat{R}}(\widehat{M}) = \widehat{R} \otimes_R \text{End}_R(M)$ . Since  $\widehat{E}/\widehat{J} = (E/J)^\wedge = E/J$ , we see that  $\text{End}_{\widehat{R}}(\widehat{M})$  is local as well, so  $\widehat{M}$  is indecomposable. Since  $M$  was arbitrary, it follows that  $R$  has bounded CM type.

For the converse we use Elkik’s theorem [E, Théorème 3] on extensions of vector bundles over Henselian pairs. Since  $\widehat{R}$  has an isolated singularity, every MCM  $\widehat{R}$ -module  $M$  is locally free on the punctured spectrum of  $\widehat{R}$ , and so by Elkik’s theorem is isomorphic to  $\widehat{N}$  for some (necessarily MCM)  $R$ -module  $N$ . It follows immediately that bounded CM type extends to  $\widehat{R}$ .  $\square$

For ascent to the Henselization we recycle an argument from [Wi1] and [LW1]. Recall [DI] that the extension  $R \rightarrow R^{\text{h}}$  is *separable*, meaning that the sequence

$$(3.1.1) \quad 0 \longrightarrow J \longrightarrow R^{\text{h}} \otimes_R R^{\text{h}} \xrightarrow{\mu} R^{\text{h}} \longrightarrow 0,$$

where  $\mu(u \otimes v) = uv$ , is split exact as a sequence of  $R^h \otimes_R R^h$ -modules. Tensoring (3.1.1) with an arbitrary finitely generated  $R^h$ -module  $N$  shows that  $N$  is a direct summand of the extended module  $R^h \otimes_R N$ , where the action of  $R^h$  on  $R^h \otimes_R N$  is by change of rings. Write  ${}_R N$  as a directed union of finitely generated  $R$ -modules  $A_\alpha$ . Then, since  $N$  is a finitely generated  $R^h$ -module,  $N$  is a direct summand of  $R^h \otimes_R A_\alpha$  for some  $\alpha$ . Thus any finitely generated  $R^h$ -module  $N$  is a direct summand of an extended module.

**Proposition 3.2.** *Let  $R$  be a CM local ring with Henselization  $R^h$ . Assume that  $R^h$  is Gorenstein on the punctured spectrum. If  $R$  has bounded CM type, then  $R^h$  does as well.*

*Proof.* Let  $M$  be an indecomposable MCM  $R^h$ -module. Put  $d = \dim(R)$ . Since  $R^h$  is Gorenstein on the punctured spectrum,  $M$  is a  $d^{\text{th}}$  syzygy of some finitely generated  $R^h$ -module  $N$ , [EG, 3.8]. By the argument above,  $N$  is a direct summand of  $R^h \otimes_R B$  for some finitely generated  $R$ -module  $B$ . Letting  $A$  be a  $d^{\text{th}}$  syzygy of  ${}_R B$ , we see (using the Krull–Schmidt Theorem over  $R^h$ ) that  $M$  is a direct summand of  $R^h \otimes_R A$ . Since  ${}_R A$  is MCM, we can write  $A$  as a direct sum of modules  $C_i$  of low multiplicity. Using Krull–Schmidt again, we deduce that  $M$  is a direct summand of some  $R^h \otimes_R C_i$ , thereby getting a bound on the multiplicity of  $M$ .  $\square$

**Theorem 3.3.** *Let  $R$  be an excellent CM local ring with at most an isolated singularity. If  $R$  has bounded CM type, then the completion  $\widehat{R}$  also has bounded CM type.*

*Proof.* As  $R$  has geometrically regular formal fibres,  $\widehat{R}$  and the Henselization  $R^h$  both also have isolated singularities [EGA, (6.5.3)]. By Proposition 3.1 and Corollary 3.2, then, bounded CM type ascends to  $R^h$  and thence to  $\widehat{R}$ .  $\square$

Theorem 3.3 allows us to verify a version of the Brauer–Thrall conjecture. The complete case of this theorem is due to Yoshino and Dieterich [Y, (6.4)].

**Theorem 3.4.** *Let  $(R, \mathfrak{m}, k)$  be an excellent equicharacteristic CM local ring with perfect residue field  $k$ . Then  $R$  has finite CM type if and only if  $R$  has bounded CM type and  $R$  has at most an isolated singularity.*

*Proof.* If  $R$  has finite CM type, then  $R$  has at most an isolated singularity by [HL], and of course  $R$  has bounded CM type. Suppose now that  $R$  has bounded CM type and at most an isolated singularity. According to Theorem 3.3,  $\widehat{R}$  also has bounded CM type. By the Brauer–Thrall theorem of Yoshino and Dieterich,  $\widehat{R}$  has finite CM type. This descends to  $R$  by [Wi1, (1.4)], and we are done.  $\square$

One cannot remove the hypothesis of excellence. For example, let  $S$  be any one-dimensional analytically ramified local domain. It is known [M, pp. 138–139] that there is a one-dimensional local domain  $R$  between  $S$  and its quotient field such that  $e(R) = 2$  and  $\widehat{R}$  is not reduced. Then  $R$  has bounded but infinite CM type by [LW2, (2.1), (0.1)], and of course  $R$  has an isolated singularity.

Proving descent of bounded CM type in general seems quite difficult. Part of the difficulty lies in the fact that, in general, there is no bound on the number of indecomposable MCM  $\widehat{R}$ -modules required to decompose the completion of an indecomposable MCM  $R$ -module. Here is an example to illustrate. Recall [RWW, Prop. 3] that when  $R$  and  $\widehat{R}$  are two-dimensional normal domains, a torsion-free  $\widehat{R}$ -module  $M$  is extended from  $R$  if and only if  $[M]$  is in the image of the natural map on divisor class groups  $\text{Cl}(R) \longrightarrow \text{Cl}(\widehat{R})$ .

**Example 3.5.** Let  $A$  be a complete local two-dimensional normal domain containing a field, and assume that the divisor class group  $\text{Cl}(A)$  has an element  $\alpha$  of infinite order. (See, for example, [Wi2, (3.4)].) By Heitmann's theorem [He], there is a unique factorization domain  $R$  contained in  $A$  such that  $\widehat{R} = A$ . Choose, for each integer  $n$ , a divisorial ideal  $I_n$  corresponding to  $n\alpha \in \text{Cl}(\widehat{R})$ . For each  $n \geq 1$ , let  $M_n := I_n \oplus N_n$ , where  $N_n$  is the direct sum of  $n$  copies of  $I_{-1}$ . Then  $M_n$  has trivial divisor class and therefore is extended from  $R$  by [RWW, Prop. 3]. However, no non-trivial proper direct summand of  $M_n$  has trivial divisor class, and it follows that  $M_n$  (a direct sum of  $n + 1$  indecomposable  $\widehat{R}$ -modules) is extended from an indecomposable MCM  $R$ -module.

It is important to note that the example above does not give a counterexample to descent of bounded CM type, but merely points out one difficulty in studying descent.

We finish with a positive result. Recall [GS] that the *Davenport constant*  $D(G)$  of a finite abelian group  $G$  is the least positive integer  $d$  such that every sequence of  $d$  (not necessarily distinct) elements of  $G$  has a non-empty subsequence whose sum is 0. It is easy to see that  $D(G) \leq |G|$ , with equality if  $G$  is cyclic.

**Proposition 3.6.** *Let  $R$  be a local normal domain of dimension two such that  $\widehat{R}$  is also a normal domain. Assume that the cokernel  $G$  of the natural map  $\text{Cl}(R) \longrightarrow \text{Cl}(\widehat{R})$  is finite. If the ranks of the indecomposable MCM  $\widehat{R}$ -modules are bounded by  $n$ , then the ranks of the indecomposable MCM  $R$ -modules are bounded by  $m := nD(G)$ . In particular, if  $\widehat{R}$  has bounded CM type, so has  $R$ .*

*Proof.* Let  $M$  be an indecomposable MCM  $R$ -module, and put  $N := \widehat{M}$ . Write

$$N \cong Z_1^{a_1} \oplus \cdots \oplus Z_k^{a_k},$$

where the  $Z_i$  are indecomposable MCM modules, each of rank at most  $n$ . It will suffice to show that  $a_1 + \cdots + a_k \leq D(G)$ .

Let  $\pi : \text{Cl}(\widehat{R}) \longrightarrow G$  be the natural map, let  $\zeta_i$  be the divisor class of  $Z_i$ , and let  $\gamma_i = \pi(\zeta_i) \in G$ . If  $a_1 + \cdots + a_k > D(G)$ , there is a sequence  $(b_1, \dots, b_k) \in \mathbb{N}^k$  such that

- (1)  $b_i \leq a_i$  for all  $i$ ;
- (2)  $b_j < a_j$  for some  $j$ ;
- (3)  $b_k > 0$  for some  $k$ ;
- (4)  $b_1\gamma_1 + \cdots + b_k\gamma_k = 0$ .

Put  $N' := Z_1^{b_1} \oplus \cdots \oplus Z_k^{b_k}$ . By (4), the divisor class of the module  $N'$  is in the image of the map  $\text{Cl}(R) \rightarrow \text{Cl}(\widehat{R})$ . Therefore  $N'$  is extended from  $R$ , by [RWW, Prop. 3], say  $N' \cong \widehat{M}'$ . Also, (1), (2) and (3) say that  $N'$  is a non-trivial proper direct summand of  $N$ . By [Wi2, (1.2)],  $M'$  is a non-trivial proper direct summand of  $M$ , contradiction.  $\square$

## REFERENCES

- [Ab] S.S. Abhyankar, *Local rings of high embedding dimension*, Amer. J. Math. **89** (1967), 1073–1077.
- [Ar] M. Arnavut, *Decomposition of modules over one-dimensional rings*, Ph.D. thesis, University of Nebraska–Lincoln, Lincoln, NE, 2002.
- [B] H. Bass, *On the ubiquity of Gorenstein rings*, Math. Z. **82** (1963), 8–28.
- [BGS] R.-O. Buchweitz, G.-M. Greuel, and F.-O. Schreyer, *Cohen–Macaulay modules on hypersurface singularities II*, Invent. Math. **88** (1987), 165–182.
- [BH] W. Bruns and J. Herzog, *Cohen–Macaulay rings*, Cambridge Stud. in Adv. Math., vol. 39, Cambridge University Press, Cambridge, 1993.
- [Ç1] N. Çimen, *One-dimensional rings of finite Cohen–Macaulay type*, Ph.D. thesis, University of Nebraska–Lincoln, Lincoln, NE, 1994.
- [Ç2] ———, *One-dimensional rings of finite Cohen–Macaulay type*, J. Pure Appl. Algebra **132** (1998), no. 3, 275–308.
- [ÇWW] N. Çimen, R. Wiegand, and S. Wiegand, *One-dimensional rings of finite Cohen–Macaulay type*, Abelian Groups and Modules (A. Facchini and C. Menini, eds.), Kluwer Academic, Dordrecht, 1995.
- [DI] F. DeMeyer and E. Ingraham, *Separable algebras over commutative rings*, Lecture Notes in Mathematics, vol. 181, Springer-Verlag, New York-Berlin, 1971.
- [D] E. Dieterich, *Reduction of isolated singularities*, Comment. Math. Helv. **62** (1987), 654–676.
- [DR] Ju.A. Drozd and A.V. Roĭter, *Commutative rings with a finite number of indecomposable integral representations*, Izv. Akad. Nauk. SSSR Ser. Mat. **31** (1967), 783–798, Russian.
- [EG] E.G. Evans and Ph. Griffith, *Szygies*, London Math. Soc. Lect. Notes Ser., vol. 106, Cambridge University Press, 1985.
- [EGA] A. Grothendieck and J. Dieudonné, *Éléments de Géométrie Algébrique IV, Partie 2*, vol. 24, Publ. Math. I.H.E.S., 1967.
- [E] R. Elkik, *Solutions d'équations à coefficients dans un anneau hensélien*, Ann. Sci. École Norm. Sup. (4) **6** (1973), 553–603.
- [GR] E. Green and I. Reiner, *Integral representations and diagrams*, Michigan Math. J. **25** (1978), 53–84.
- [GS] A. Geroldinger and R. Schneider, *On Davenport's Constant*, J. Comb. Theory Ser. A, **61** (1992), 147–152.
- [He] R. Heitmann, *Characterizations of completions of unique factorization domains*, Trans. Amer. Math. Soc. **337** (1993), 379–387.
- [HL] C. Huneke and G. Leuschke *Two theorems about maximal Cohen–Macaulay modules*, Math. Ann. **324** (2002), 391–404.
- [LW1] G. Leuschke and R. Wiegand, *Ascent of finite Cohen–Macaulay type*, J. Algebra **228** (2000), 674–681.
- [LW2] ———, *Hypersurfaces of bounded Cohen–Macaulay type*, J. Pure Appl. Algebra (to appear).
- [M] E. Matlis, *One-dimensional Cohen–Macaulay Rings*, Lect. Notes in Math., vol. 52, Springer-Verlag, Berlin, 1973.
- [R] D.E. Rush, *Rings with two-generated ideals*, J. Pure Appl. Algebra **73** (1991), no. 3, 257–275.
- [RWW] C. Rotthaus, D. Weston, and R. Wiegand, *Indecomposable Gorenstein modules of odd rank*, J. Algebra **214** (1999), 122–127.

- [S] J.D. Sally, *Numbers of generators of ideals in local rings*, Marcel Dekker Inc., New York, 1978.
- [SS] G. Scheja and U. Storch, *Differentielle Eigenschaften der Localisierungen analytischer Algebren*, Math. Ann. **197** (1972), 137–170.
- [SV] J.D. Sally and W. Vasconcelos, *Stable rings*, J. Pure Appl. Algebra **4** (1974), 319–336.
- [We] D. Weston, *On descent in dimension two and non-split Gorenstein modules*, J. Algebra **118** (1988), 263–275.
- [Wi1] R. Wiegand, *Local rings of finite Cohen–Macaulay type*, J. Algebra **203** (1998), 156–168.
- [Wi2] ———, *Direct-sum decompositions over local rings*, J. Algebra **240** (2001), 83–97.
- [Y] Y. Yoshino, *Cohen–Macaulay modules over Cohen–Macaulay rings*, London Math. Soc. Lect. Notes Ser., vol. 146, Cambridge University Press, 1990.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF KANSAS, LAWRENCE, KS 66045

*E-mail address:* [gleuschke@math.ukans.edu](mailto:gleuschke@math.ukans.edu)

*URL:* <http://www.leuschke.org>

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NEBRASKA–LINCOLN, LINCOLN, NE 68588-0323

*E-mail address:* [rwiegand@math.unl.edu](mailto:rwiegand@math.unl.edu)

*URL:* <http://www.math.unl.edu/~rwiegand>